Communicating Difficult News

Adamowski K, Dickinson G, Weitzman B, Roessler C, Carter-Snell C. Sudden unexpected death in the emergency department: caring for the survivors. *CMAJ*.1993; 149(10):1445-1451.

Baile WF, Buckman R, Lenzi R, Glober G, Beale EA, Kudelka AP. SPIKES-A six-step protocol for delivering bad news: application to the patient with cancer. *Oncologist*. 2000; 5(4):302-311.

Barclay JS, Blackhall LJ, Tulsky JA. Communication strategies and cultural issues in the delivery of bad news. *J Palliat Med*. 2007; 10(4):958-797.

Benenson RS, Pollack ML. Evaluation of emergency medicine resident death notification skills by direct observation. *Acad Emerg Med*. 2003;10(3):219–223.

Bruera E, Palmer JL, Pace E, et al. A randomized, controlled trial of physician postures when breaking bad news to cancer patients. *Palliat Med*.2007; 21(6):501-505.

Buckman R. *How to Break Bad News: A guide for Health Care Professions*. Baltimore, MD: Johns Hopkins University Press; 1992.

Fallowfield L, Jenkins V. Communicating sad, bad, and difficult news in medicine. *Lancet*. 2004; 363(9405):312-319.

Hobgood C, Harward D, Newton K, Davis W. The educational intervention

"GRIEV_ING" improves the death notification skills of residents. *Acad Emerg Med.* 2005; 12(4):296-301.

Iserson K. *Grave Words: Notifying Survivors about Sudden Unexpected Deaths*. Tucson, AZ: Galen Press; 1999.

Knopp R, Rosenzweig S, Bernstein E, Totten V. Physician–patient communication in the emergency department, part 1. *Acad Emerg Med.* 1996; 3(11):1065–1069.

McCaig LF, Burt CW. National Hospital Ambulatory Medical Care Survey: 2002

Emergency Department Summary. Advance data from vital and health statistics; no 340. Hyattsville, Maryland: National Center for Health Statistics. 2004.

http://www.cdc.gov/nchs/data/ad/ad340.pdf. Accessed June 18, 2008.

Parrish GA, Holdren KS, Skiendzielewski JJ, Lumpkin OA. Emergency department experience with sudden death: a survey of survivors. *Ann Emerg Med.* 1987; 6(7):792–796.

Physician–patient communication in the emergency department. Part 2: Communication strategies for specific situations. SAEM Task Force on Physician–Patient

Communication. Acad Emerg Med. 1996;3(12):1146–1153.

Quest TE, Otsuki JA, Banja J, Ratcliff JJ, Heron SL, Kaslow NJ. The use of standardized patients within a procedural competency model to teach death disclosure. *Acad Emerg Med.* 2002; 9(11):1326–1333.

Spinetta JJ, Maloney LJ. The child with cancer: patterns of communication and denial. *J Consult Clin Psychol*. 1978;46(6):1540–1541.

Suchman AL, Markakis K, Beckman HB, Frankel R. A model of empathic communication in the medical interview. *JAMA*. 1997;277(8):678–682.

Swisher LA, Nieman LZ, Nilsen GJ, Spivey WH. Death notification in the emergency department: a survey of residents and attending physicians. *Ann Emerg Med*. 1993;22(8):1319–1323.

Sykes N. Medical students' fears about breaking bad news. *Lancet*. 1989;2(8662):564. Tolle SW, Cooney TG, Hickam DH. A program to teach residents humanistic skills for notifying survivors of a patient's death. *Acad Med*. 1989;64(9):505–506.

United States Department of Health and Human Services. Office for Civil Rights. Privacy Rule Summary. http://www.hhs.gov/ocr/privacysummary.pdf Accessed June 17, 2008.

Vaidya VU, Greenberg LW, Patel KM, Strauss LH, Pollack MM. Teaching physicians how to break bad news: a 1-day workshop using standardized parents. *Arch Pediatr Adolesc Med.* 1999;153(4):419–422.

Walters DT, Tupin JP. Family grief in the emergency department. *Emerg Med Clin N Am.* 1991;9(1):189–206.

Wolraich ML, Reiter S. Training physicians in communication skills. *Dev Med Child Neurol*. 1979;21(6):773–778.